

Medical treatment options for women with Heavy Menstrual Bleeding (HMB) due to fibroids (HMB-L)

Information for patients and carers

Does this information apply to me?

Yes, if you are a woman who has heavy periods that are associated with fibroids.

It does not cover:

- heavy periods caused by endometriosis, hormone therapy or bleeding disorders
- bleeding that is not related to your menstrual cycle.

Are heavy periods disrupting your life?

Every woman is different and the amount of blood each woman loses during her period varies widely from one person to another. If heavy periods are disrupting your life, your doctor should be able to offer treatments to help.

Working with you

Your doctor should talk with you about heavy periods. They should explain any tests and treatments so that you can decide together what is best for you. Your family or carer can be involved in helping to make decisions, but only if you agree.

Some treatments or care described here may not be suitable for you. If you think that your treatment does not match this advice, talk to your care team.

Your doctor will be able to offer a number of different drug treatments to help you. Some of the treatments are also contraceptives. Your doctor should discuss the benefits and risks of each treatment with you. If one treatment isn't suitable for you, or if you try one treatment and it doesn't work, it may be possible to try another option. Some of the treatments make your periods lighter and some may stop the bleeding completely. You should be given information explaining the different options, and be allowed time to make your decision.

Treatment options

Tranexamic and/or Mefanamic Acid

What are they? Non-hormonal tablets taken from the start of your period for up to 4/5 days.

How do they work? Tranexamic acid helps the blood in the womb to form clots, which reduces the amount of bleeding. Mefanamic acid is an anti-inflammatory medication.

Impact on fertility: none.

Possible unwanted side effects: may cause indigestion and/or diarrhoea. Women who have had a blood clot in their leg or lung should not use tranexamic acid.

Ulipristal Acetate

What is it? Tablets that are taken once daily for 3 months. You can have up to 4 courses of treatment of 3 months each, waiting for withdrawal bleeding and a period before starting your next course of tablets.

How does it work? It acts on the progesterone receptor.

Impact on fertility: reduces fertility while taking the tablets but it is not a contraceptive in the regime prescribed. It's effects are reversible.

Possible unwanted side effects: Many women stop having periods when taking this drug. Common unwanted effects include stomach pain, feeling sick, headaches, dizziness, tiredness, weight gain, hot flushes and breast tenderness.

Oral Progestogen (Norethisterone)

What is it? Tablets taken 2 to 3 times a day from the 5th to the 26th day of your cycle (counting the first day of your period as day 1).

How does it work? Prevents the lining of the womb from growing quickly.

Version: 0000 Approved: 0000 Review: 0000

Approved by: Clinical Policy, Documentation & Information Group

Impact of fertility: not an effective contraceptive but will decrease the chance of becoming pregnant. Fertility potential will return on stopping the medication.

Possible unwanted side effects: Common: weight gain; bloating; breast tenderness; headaches; acne (usually minor and short lived).

Injected or Implanted Progestogen

What is it? An injection of progestogen (similar to the natural hormone, progesterone). An implant is also available that releases progestogen slowly for up to 3 years.

How does it work? Prevents the lining of the womb from growing quickly.

Impact of fertility: Fertility returns on stopping the medication although time-frame of return of fertility will vary depending upon preparation administered. The progestogen injection can rarely cause a delay in fertility of up to 18 months.

Possible unwanted side effects: Common: weight gain; irregular bleeding; absence of periods; premenstrual symptoms (including bloating, fluid retention, breast tenderness). Less common: bone density loss.

Levonorgestrel- releasing Intrauterine System (IUS; e.g. Mirena®)

What is it: A small plastic device (intra-uterine system; IUS) that is placed in the womb and slowly releases progestogen (which acts like the natural hormone, progesterone). It can remain in the womb for up to 5 years (or longer if you are over 45 when it is inserted). Sometimes it is necessary to have this device inserted at the same time as a hysteroscopy (a small camera inserted into the womb), particularly if your fibroids are large. Not all womb cavities are suitable; if the fibroids distort the cavity significantly you may not be suitable for this treatment option. Your doctor will advise you on this.

How does it work: Prevents the lining of the womb from growing quickly. Approximately 4 in 5 women will have a significant reduction in their menstrual blood loss (up to 90% reduction).

Impact on fertility: it is a contraceptive but fertility returns within a month once it is removed.

Possible complications/unwanted side effects: Commonly causes irregular bleeding or spotting that may last for over 6 months; breast tenderness, acne or headaches may occur but are generally minor and short lived. Less commonly it will stop periods completely. Sometimes this device can fall out, particularly if your fibroids are large. Occasionally the IUS can perforate the womb when it is inserted and rarely women require a keyhole procedure for this complication. The risk of infection is increased for 2-3 weeks after insertion and a course of antibiotics may be required.

Gonadotrophin- releasing Hormone Analogue e.g. Decapeptyl/Lupron/Prostap

What is it? An injection that stops the body producing the hormones oestrogen and progesterone, effectively creating a temporary (reversible) menopause.

How does it work? Prevents the menstrual cycle. Reduces fibroid size and volume. It can be used for 6 months on its own or for longer with hormone replacement therapy (up to 2 years). This option is not normally suitable as a long term treatment.

Impact of fertility: Non-hormonal contraception is advised while using this option. It is unlikely that you will be able to conceive on this treatment but fertility returns rapidly on stopping.

Possible unwanted side effects: Common: menopause- like symptoms (for example, hot flushes, increased sweating, vaginal dryness). Less common: risk of osteoporosis with longer-term use. Your doctor may offer you HRT to take alongside this treatment to prevent/alleviate these effects.

Note: The most common unwanted effects may be experienced by 1 in 100 women. Less common unwanted effects are those experienced by 1 in 1000 women. Rare unwanted effects are not shown here.

Further Information

Contact Telephone Numbers:	Edinburgh Royal Infirmary	0131 242 2551
	Chalmers Centre	0131 526 1070
	St John's Hospital	01506 524383

Information in these leaflets has been adapted from the National Institute for Health and Care Excellence (<https://www.nice.org.uk/>) and Royal College of Obstetricians and Gynaecologists (<https://www.rcog.org.uk/en/guidelines-research-services/>) guidelines.

Version: 0000 Approved: 0000 Review: 0000

Approved by: Clinical Policy, Documentation & Information Group